

A.

8.10 The sample space of all possible sequences is {HHH, HHT, HTH, THH, HTT, THT, TTH, TTT}

a. For each sequence in the sample space, count the number of tails. Then count how many outcomes there are for each possible number of tails. Divide by 8 because all outcomes in the sample space are equally likely. The distribution (*pdf*) is:

k	0	1	2	3
$P(X=k)$	1/8	3/8	3/8	1/8

b. For each sequence in the sample space, take the difference (Y) between the number of heads and the number of tails. For instance, for HHT, $Y = 2$ (heads) $- 1$ (tails) $= 1$. List all possible differences, count how many sequences in the sample space have each possible value, then divide by 8 to get the probability of that value. The distribution (*pdf*) is:

k	-3	-1	1	3
$P(Y=k)$	1/8	3/8	3/8	1/8

c. The sum of heads and tails is 3 for every sequence which means $P(Z=3) = 1$ so the distribution (*pdf*) is:

k	3
$P(Z=k)$	1

8.21 The distribution of the sum is given in Example 8.5. The expected value is

$$\mu = E(X) = 2 \times \frac{1}{36} + 3 \times \frac{2}{36} + 4 \times \frac{3}{36} + \dots + 10 \times \frac{3}{36} + 11 \times \frac{2}{36} + 12 \times \frac{1}{36} = \frac{252}{36} = 7$$

8.22 $\sigma = 2.415$. The formula is $\sigma = \sqrt{\sum (x - \mu)^2 p(x)}$ where the sum is over all values of x . Here, the mean is $\mu = 7$ (see Exercise 8.10). Details of the calculations are:


X	2	3	4	5	6	7	8	9	10	11	12
$x - \mu$	-5	-4	-3	-2	-1	0	1	2	3	4	5
$(x - \mu)^2$	25	16	9	4	1	0	1	4	9	16	25
$p(x)$	1/36	2/36	3/36	4/36	5/36	6/36	5/36	4/36	3/36	2/36	1/36

$$\begin{aligned} \sigma &= \sqrt{25 \frac{1}{36} + 16 \frac{2}{36} + 9 \frac{3}{36} + 4 \frac{4}{36} + 1 \frac{5}{36} + 0 \frac{6}{36} + 1 \frac{5}{36} + 4 \frac{4}{36} + 9 \frac{3}{36} + 16 \frac{2}{36} + 25 \frac{1}{36}} \\ &= \sqrt{210/36} = 2.415. \end{aligned}$$

- 8.36** Use a binomial distribution with $n = 10$, $p = .5$. Let X = number of games won by human. The answers can be found using any of the methods discussed in Section 8.4, including the use of Minitab or Excel.
- $P(X = 5) = .2461$.
 - $P(X = 3) = .1172$. If computer wins 7 games, then human wins 3 games.
 - $P(X \geq 7) = .1719$, calculated as $P(X = 7) + P(X = 8) + P(X = 9) + P(X = 10) = .1172 + .0439 + .0098 + .0010 = .1719$. Equivalently, $P(X \geq 7) = 1 - P(X \leq 6) = 1 - .8281$.

- 8.46**
- X is a uniform random variable (and it is continuous).
 - X ranges from 0 to 100 and the area under any density curve is 1, so $f(x) = 1/100 = .01$ for all x between 0 and 100. This creates a rectangle (with area=1) similar to Figure 8.2. *Note:* $f(x) = 0$ for any x not between 0 and 100.
 - $P(X \leq 15 \text{ seconds})$ is the area of the rectangle from 0 to 15 seconds. The interval width is 15 and the height is $1/100$, so the answer is $(15)(1/100) = .15$.
 - $P(X \geq 40 \text{ seconds})$ is the area of the rectangle between 40 and 100. The interval width is 60 and the height is $1/100$ so the answer is $(60)(1/100) = .60$.

e.



f. The expected value or mean is 50. The distribution is symmetric, so the mean equals the median. For a uniform random variable, the median is at the middle of the range of possible values.

- 8.78** Solve this problem using a binomial distribution with $n=10$ and $p=0.1$. $P(X \geq 3) = 1 - P(X \leq 2) = 1 - .9298 = .0702$. *Note:* If using the binomial formula to find $P(X \leq 2)$ by hand, calculate $P(X=0) + P(X=1) + P(X=2)$ which will be $.3487 + .3874 + .1937 = .9298$.

B. Suppose that a random variable X has cdf given by the function $F(k) = \sqrt{k}$ for values of k between 0 and 1. (And $F(k)=0$ for negative k, and $F(k)=1$ for k greater than 1.) Find the expectation and standard deviation of X. Show all your work.

$$\text{CDF is } F(k) = \begin{cases} \sqrt{k} & 0 \leq k \leq 1 \\ 0 & k < 0 \\ 1 & k > 1 \end{cases} \quad \text{To find the expectation of X, } E(X), \text{ we need to define the pdf of X, } f(k).$$

$$f(k) \text{ is the derivative of } F(k). \text{ So } f(k) = \begin{cases} \frac{1}{2}k^{-1/2} & 0 \leq k \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$\text{Now we can go ahead and find } E(X). E(X) = \int_0^1 \frac{1}{2}k^{-1/2} * k \, dk = \int_0^1 \frac{1}{2}k^{1/2} \, dk = \frac{1}{3}k^{3/2} \Big|_0^1 = \frac{1}{3} - 0 = \frac{1}{3}$$

To find the standard deviation of X, we need to find $E(X^2)$ first.

$$E(X^2) = \int_0^1 \frac{1}{2}k^{-1/2} * k^2 \, dk = \int_0^1 \frac{1}{2}k^{3/2} \, dk = \frac{1}{5}k^{5/2} \Big|_0^1 = \frac{1}{5} - 0 = \frac{1}{5}$$

$$\text{The standard deviation of X is the square root of its variance. } \text{Var}(X) = E(X^2) - (E(X))^2 = \frac{1}{5} - \left(\frac{1}{3}\right)^2 = \frac{4}{45}$$

$$\text{The standard deviation of X} = \sqrt{\frac{4}{45}} \approx 0.2981$$

C. Suppose that a continuous random variable Y has pdf that is equal to $f(k) = c(k)(k-1)$ for some constant c and for all k in the interval (1,3). Assume that the pdf is zero outside the interval (1,3). First, find the value of c. Then, calculate $E(Y)$. Show all your work.

$$f(k) = \begin{cases} c(k)(k-1) & 1 \leq k \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

We can solve for c since we know that every pdf has to integrate to 1.

$$\text{So } 1 = \int_1^3 c k (k-1) \, dk = c \int_1^3 (k^2 - k) \, dk = c \left(\frac{k^3}{3} - \frac{k^2}{2} \Big|_1^3 \right) = c \left(\frac{26}{3} - \frac{8}{2} \right) = c \frac{14}{3} \rightarrow c = \frac{3}{14}$$

$$E(Y) = \frac{3}{14} \int_1^3 k * (k^2 - k) \, dk = \frac{3}{14} \int_1^3 (k^3 - k^2) \, dk = \frac{3}{14} \left(\frac{k^4}{4} - \frac{k^3}{3} \Big|_1^3 \right) = \frac{3}{14} \left(\frac{80}{4} - \frac{26}{3} \right) = \frac{17}{7} \approx 2.4286$$